

2
3 **Population consequences of migratory variability differ between flyways**

4 Robert Patchett¹, Tom Finch^{2,3} and Will Cresswell^{1,*}

5 ¹Centre for Biological Diversity, University of St Andrews, St Andrews, Fife KY16 9TH, UK

6 ²RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, The Lodge, Sandy
7 SG19 2DL, UK

8 ³Department of Zoology, University of Cambridge, Downing St, Cambridge CB2 3EJ, UK

9 *Lead Contact: wrlc@st-and.ac.uk , +44 1334 463010

10
11 eTOC

12 Patchett et al. show that climate change, habitat loss and migration distance may influence population
13 trends in migrant landbirds, but their relative effect varies between two major flyways.

14
15 Many populations of long distance migratory birds are declining, with candidate drivers including
16 climate change shifting the location of suitable non-breeding habitat, outright anthropogenic habitat
17 loss, or a combination of both [1]. The size of the non-breeding area over which a population spreads
18 (migratory spread), and migration distance (or number of stop-over sites), however, may determine a
19 species' susceptibility to such environmental change. Low migratory spread is predicted to increase
20 vulnerability to climate-driven shifts in the location of suitable non-breeding habitat, because
21 individuals are less likely to encounter newly suitable sites. All else being equal, outright loss of non-
22 breeding habitat should result in population declines regardless of migratory spread. However, if
23 migratory spread is low and habitat loss is uneven we would expect to sample some species which
24 are experiencing habitat loss and some species which are not. In contrast, under high spread (or
25 when habitat loss is evenly distributed), any sampled species is likely to experience habitat loss
26 (Figure 1C). As migration distance increases so the probability of encountering a stop-over site with
27 negative environmental change increases [3] (Figure 1D). Therefore, the presence of a positive

relationship between species' population trends and migratory spread or a negative relationship between species' population trends and migration distance, should indicate whether climate-change-driven shifts in non-breeding habitat location, or environmental change at stopover sites, respectively, are likely drivers of migratory declines. These relationships may vary between flyways, which differ profoundly in their geography, climate variation and human population change. Here we show that migratory connectivity is correlated with migrant population trends and that low spread is associated with declining populations in the Neotropic flyway, but the reverse occurs in the Afro-Palearctic flyway. Thus climate change may be more important in Neotropic migrant population declines whereas habitat loss may be more important in the Afro-Palearctic.

We tested for relationships between species' population trends and their mean spread on the non-breeding ground, as well as their migration distance, and whether these relationships varied between the Neotropic and Afro-Palearctic flyways. We used all available published individual migration tracks (tracked from breeding to non-breeding grounds) of 875 adult landbirds from 122 populations of 48 species (Figure 1A and B) to calculate migratory spread (the mean inter-individual distance between birds from the same population on the non-breeding ground; data as in Finch et al. 2017 [4], with the addition of 17 more recent studies). We modelled each species' population status, using the most current IUCN assessment of declining, stable and increasing populations [5], against its non-breeding ground mean spread and mean distance of migration. We also included whether the species was in the Neotropic or Afro-Palearctic flyway, and interactions dependent on flyway, and controlled for the latitudinal zone of wintering (approximately the Caribbean and Central America versus South America, and Africa north and south of the equator), the longitude and latitude of the breeding population, and the phylogeny of the migratory species considered [4].

Scoring declining, stable and increasing populations as -1, 0 and 1, respectively, the mean population trend was -0.2 (47% of species declining, 27% increasing) in the Neotropic flyway and -0.55 (64% of species declining, 9% increasing) in the Afro-Palearctic flyway. In the Neotropic flyway, species with low spread (high connectivity) were significantly more likely to have declining populations ($t_{11.8} = 3.7$, $P < 0.01$), whilst in the Afro-Palearctic flyway, species with high spread (low connectivity) were more likely to have declining populations ($t_{27.7} = -2.2$ $P = 0.033$; interaction between flyway and migratory spread, $t_{41.3} = 3.9$, $P < 0.001$; Table S2, Figure 1G). In the Neotropic flyway, species with greater migratory distances were significantly more likely to have declining populations ($t_{9.1} = -3.5$, $P < 0.01$),

whilst in the Afro-Palearctic flyway there was no relationship ($t_{27.4} = -0.2$; interaction between flyway and migration distance, $t_{39.6} = -2.2$, $P=0.034$; Table S2, Figure 1H). The fixed effects of flyway, migration distance and spread (and interactions between flyway and migration distance or spread), accounted for 30% of the variation in population trend, with phylogeny contributing an additional 5%. Migratory spread and migration distance accounted for 55% and 13% of variation in population trends in the Neotropic and Afro-Palearctic flyways respectively.

The positive correlation between population trend and migratory spread suggests that climate-change-driven shifts in the location of non-breeding habitat could be a driver of population declines in the Neotropic flyway. In contrast, there was a weak negative relationship between population trend and migratory spread in the Afro-Palearctic flyway, suggesting that habitat loss is occurring unevenly across Africa with a diluted impact on species with low spread. Long-term climatic conditions in South America may have been more stable in comparison to Africa [7], where climate varies across a continental scale between decades [6]. Migrant birds in the Afro-Palearctic flyway may then have already evolved generalist traits and a bet-hedging strategy of wide migratory spread in response [2]. Although habitat loss through land use change occurs in both flyways, human population growth (a reasonable proxy for rates of anthropogenic habitat transformation) has, and still is, occurring at a much greater rate in Africa but it is uneven and concentrated regionally (i.e. countries including and around Nigeria, and Uganda) [8]. The savannahs of sub-Saharan Africa, where most Afro-Palearctic species spend the non-breeding season, also have fewer protected areas than African tropical forests [9]. The differences between flyways in the effects of migration distance may then reflect current climate change and habitat loss in Central America and the Caribbean removing stop-over sites now, whereas at equivalent latitudes in Africa these were removed five thousand years ago with the formation of the Sahara [10].

Other factors acting on breeding and non-breeding grounds are also likely to be important drivers such as phenology mismatch [3]. Nevertheless, our results highlight how non-breeding migratory spread (migratory connectivity) might help explain general population trends of long-distance migrant birds, because migratory spread is likely to be an indication of a suite of bet-hedging adaptations to past variable climate and so a guide to future climate change responses, and how these might differ across flyways.

88 Author Contributions

89 All authors contributed equally to conceptualization, data collation, analysis and writing the paper.

90

91 Acknowledgments

92 We are extremely grateful to and thank all of the researchers that have contributed to and published
93 migratory bird movement data. We thank Birdlife International and their numerous contributors that
94 enable them provide population status data. We thank Christian Rutz and two anonymous referees
95 for helpful comments and suggestions.

96

97 Supplemental Information

98 Document S1. Experimental Procedures and Two Tables

99

100 Declaration of Interests

101 The authors declare no competing interests.

102

103 References

- 104 1. Vickery, J.A., Ewing, S., Smith, K.W., Pain, D.J., Bairlein, F., Škorpilová, J., and Gregory, R.D.
105 (2014). The decline of Afro-Palearctic migrants and an assessment of potential causes. *Ibis*.
106 156, 1–22.
- 107 2. Cresswell, W. (2014). Migratory connectivity of Palaearctic-African migratory birds and their
108 responses to environmental change: The serial residency hypothesis. *Ibis*. 156, 493–510.
- 109 3. Jones, T., and Cresswell, W. (2010). The phenology mismatch hypothesis: are declines of
110 migrant birds linked to uneven global climate change? *J. Anim. Ecol.* 79, 98–108.
- 111 4. Finch, T., Butler, S., Franco, A., and Cresswell, W. (2017). Low migratory connectivity is
112 common in long-distance migrant birds. *J. Anim. Ecol.* 38, 42–49.
- 113 5. BirdLife International (2017). IUCN Red List for Birds. Available at: <http://www.birdlife.org>
114 [Accessed October 2, 2017].


- 115 6. Nicholson, S.E. (2001). Climatic and environmental change in Africa during the last two
116 centuries. *Clim. Res.* 17, 123–144.
- 117 7. Anhuf, D., Ledru, M.P., Behling, H., Da Cruz, F.W., Cordeiro, R.C., Van der Hammen, T.,
118 Karmann, I., Marengo, J.A., De Oliveira, P.E., Pessenda, L., *et al.* (2006). Paleo-environmental
119 change in Amazonian and African rainforest during the LGM. *Palaeogeogr. Palaeoclimatol.*
120 *Palaeoecol.* 239, 510–527.
- 121 8. Lutz, W., and Samir, K.C. (2010). Dimensions of global population projections: what do we
122 know about future population trends and structures? *Philos. Trans. R. Soc. B Biol. Sci.* 365,
123 2779–2791.
- 124 9. Aleman, J.C., Blarquez, O., and Staver, C.A. (2016). Land-use change outweighs projected
125 effects of changing rainfall on tree cover in sub-Saharan Africa. *Glob. Chang. Biol.* 22, 3013–
126 3025.
- 127 10. Foley, J.A., Coe, M.T., Scheffer, M., and Wang, G. (2003). Regime Shifts in the Sahara and
128 Sahel: Interactions between Ecological and Climatic Systems in Northern Africa. *Ecosystems*
129 6, 524–539.

Figure 1.


Climate change, habitat loss and migration distance influence population declines in migrant landbirds, but their effect varies between two major flyways.

(A) Mercator projection of the Neotropic and (B) Afro-Palearctic migration systems showing breeding and non-breeding locations of all individuals used in analysis. (C) Representation of how species with low and high non-breeding ground spread (a measure of migratory connectivity) are predicted to be affected under scenarios of climate change shifting habitat or outright habitat loss (either evenly across range or uneven and localised into one main area) between two time periods. With climate change, high spread species are already spreading into shifted habitats as part of their bet-hedging strategy so there is no population change, whereas low spread species may now miss the shifted target habitat and show population declines. With outright habitat loss, however, all species are predicted to suffer declines, but if habitat loss is not even then habitat loss anywhere will always affect a high non-breeding spread population, but only those low non-breeding spread populations that have a range that encompasses the area of habitat change. Uneven habitat loss will therefore affect more high than low spread species. (D) Representation of how the number of stop-overs will increase with migration distance. As the number of stop-overs increases, so the probability of encountering the negative effects of climate change or habitat loss at any one of these stop-overs increases leading to population declines. (E) Predicted population trends in relation to non-breeding ground spread for climate change (P1) and habitat loss (even P2E and uneven P2U), and (F) in relation to migration distance (P3). (G) Observed relationships of population trend with non-breeding ground spread, and (H) migration distance. Lines in (G) and (H) are plotted from GLMMs \pm 1SE; line colour corresponds to predictions in (E) and (F).


Data




Hypotheses



Predictions



Results



Supplementary Information: Population consequences of migratory variability differ between flyways

Robert Patchett, Tom Finch and Will Cresswell

SUPPLEMENTAL EXPERIMENTAL PROCEDURES

Migration Data

We used an updated version of the database in Finch *et al.* [S1] that contained breeding and wintering locations for 875 individuals, from 122 populations of 48 species (see Table S1 and Supplementary References for all studies used in analysis). Data came from a comprehensive search of peer-reviewed tracking studies for all European and North American species classed (according to BirdLife; <http://www.birdlife.org/datazone/species/search>) as migratory land-birds by entering the terms [latin name] AND migra* AND (gps OR geolo* OR satellite) into the Web of Science online library. From these studies, breeding and non-breeding (i.e. the site where an individual spent the majority of the wintering, non-breeding period after migration) locations of individual birds were extracted (or approximated from plotted map locations using Google Earth when precise coordinates were not given).

Our data are prone to two potential sources of error; imprecision in the translation of data from published figures to latitude-longitude coordinates via Google Earth ('translation error'), and inaccuracy of solar geolocator-derived positions in the original published data ('geolocator error'). The sensitivity of our results to these sources of error was fully explored in Finch *et al.* [S1]. Results were little affected suggesting that errors were unbiased, and effects were relatively small and so are not considered further.

As in Finch *et al.* [S1], we restricted our study to the autumn (fall) migration of adult birds that were tagged on the breeding ground; we removed species with a sample size of one; we defined the Afro-Palearctic system as populations breeding in Europe west of 65°E that have a non-breeding area in Africa south of 20°N; we defined the Neotropic system as all populations breeding in North America with a non-breeding area south of 30°N; we split each systems' non-breeding area into northern and southern zones based on the distinct change in land to sea ratio (split at 12°N in the Neotropic and 4°N in the Afro-Palearctic system); and we classed birds tagged within 100 km of each other as

coming from the same population. Where there was more than one non-breeding site reported for an individual, we selected the non-breeding location based on length of stay (i.e. identifying the main wintering area), unless that information was unavailable when we selected the first reported non-breeding site. Of the 875 individual birds used for analysis, only 40 were reported to have wintered in more than one location.

We used the mean pairwise great circle distance between non-breeding locations for individuals from the same population as a measure of population spread (as in Finch *et al.* [S1]). We then calculated mean species-level spread (referred to as *spread*) from these measures of mean spread, averaged across all populations sampled for each species.

We used the mean great circle distance between the breeding and non-breeding locations for individuals from the same population to calculate a measure of migration distance (as in Finch *et al.* [S1]). We calculated mean species-level migration distance from these measures of migration distance, averaged across all populations sampled for each species.

Population Trend

We used species-level population trend data (*increasing, stable, decreasing* or *unknown*) published online by BirdLife International [S2] to classify a trend for each species where we could calculate mean species non-breeding spread (Table S1). Species with *unknown* trend were omitted from analysis.

Statistical Analysis

We used GLMMs to explore the extent to which population trend (*increasing, stable, decreasing* coded as 1, 0, -1 respectively) depends on spread and migration distance, with the total sample size reflecting the number of each different species in the two flyways (N = 48). We included fixed effects for spread (we transformed spread using the natural logarithm to ensure normally distributed residuals), migration distance (great circle distance between mean breeding and non-breeding sites), breeding longitude and latitude (centred for each system), zone and system. Continuous predictor variables were centred and scaled. We also included interactions between system and migration distance, and system and species spread. To test for higher level effects of taxonomy we included a random effect of family nested within order. Models were fitted using the Lme4 package in R. Model fit was assessed by visual inspection of residuals plotted against fitted values and quantile plots. Full

212 models including all terms of interest and confounding variables and minimum adequate models
213 including only significant terms were compared and gave similar results (Table S2).

214

215 Table S1. All species used in analysis with respective population trend from BirdLife International
 216 [S2].

Species (common name)	Family	Order	Population trend	System	References
Alpine swift	Apodidae	Apodiformes	stable	Afro-Palearctic	[S3]
Aquatic warbler	Acrocephalidae	Passeriformes	decreasing	Afro-Palearctic	[S4]
Barn swallow	Hirundinidae	Passeriformes	decreasing	Afro-Palearctic	[S5]
Common cuckoo	Cuculidae	Cuculiformes	decreasing	Afro-Palearctic	[S6,S7]
Common nightingale	Muscicapidae	Passeriformes	stable	Afro-Palearctic	[S8]
Common nightjar	Caprimulgidae	Caprimulgiformes	decreasing	Afro-Palearctic	[S9–S11]
Common redstart	Muscicapidae	Passeriformes	increasing	Afro-Palearctic	[S12]
Common swift	Apodidae	Apodiformes	stable	Afro-Palearctic	[S13–S16]
Cyprus wheatear	Turdidae	Passeriformes	stable	Afro-Palearctic	[S17]
Egyptian vulture	Accipitridae	Accipitriformes	decreasing	Afro-Palearctic	[S18,S19]
Eleonora’s falcon	Falconidae	Accipitriformes	increasing	Afro-Palearctic	[S20–S22]
European hoopoe	Upupidae	Bucerotiformes	decreasing	Afro-Palearctic	[S23,S24]
European roller	Coraciidae	Coraciiformes	decreasing	Afro-Palearctic	[S25]
Great reed warbler	Acrocephalidae	Passeriformes	decreasing	Afro-Palearctic	[S26,S27]
Hobby	Falconidae	Accipitriformes	decreasing	Afro-Palearctic	[S28,S29]
Honey buzzard	Accipitridae	Accipitriformes	decreasing	Afro-Palearctic	[S30]
House martin	Hirundinidae	Passeriformes	decreasing	Afro-Palearctic	[S31]
Lesser kestrel	Falconidae	Accipitriformes	stable	Afro-Palearctic	[S32,S33]
Lesser spotted eagle	Accipitridae	Accipitriformes	stable	Afro-Palearctic	[S34]
Marsh harrier	Accipitridae	Accipitriformes	increasing	Afro-Palearctic	[S35]
Montagu’s harrier	Accipitridae	Accipitriformes	decreasing	Afro-Palearctic	[S36–S38]
Northern wheatear	Muscicapidae	Passeriformes	decreasing	Afro-Palearctic	[S39,S40]
Ortolan bunting	Emberizidae	Passeriformes	decreasing	Afro-Palearctic	[S41,S42]
Pallid harrier	Accipitridae	Accipitriformes	decreasing	Afro-Palearctic	[S43]
Pied flycatcher	Muscicapidae	Passeriformes	decreasing	Afro-Palearctic	[S44]
Red-backed shrike	Laniidae	Passeriformes	decreasing	Afro-Palearctic	[S45,S46]
Sand martin	Hirundinidae	Passeriformes	decreasing	Afro-Palearctic	[S31]
Semi-collared flycatcher	Muscicapidae	Passeriformes	decreasing	Afro-Palearctic	[S47]
Short-toed eagle	Accipitridae	Accipitriformes	stable	Afro-Palearctic	[S48,S49]
Tawny pipit	Motacillidae	Passeriformes	stable	Afro-Palearctic	[S50]
Thrush nightingale	Muscicapidae	Passeriformes	stable	Afro-Palearctic	[S51]
Turtle dove	Columbidae	Columbiformes	decreasing	Afro-Palearctic	[S52,S53]
Willow warbler	Phylloscopidae	Passeriformes	decreasing	Afro-Palearctic	[S54]
Bullocks oriole	Icteridae	Passeriformes	stable	Neotropic	[S55]
Blackpoll warbler	Parulidae	Passeriformes	decreasing	Neotropic	[S56]
Bobolink	Icteridae	Passeriformes	decreasing	Neotropic	[S57]
Broad-winged hawk	Accipitridae	Accipitriformes	increasing	Neotropic	[S58]
Eastern kingbird	Tyrannidae	Passeriformes	decreasing	Neotropic	[S59]
Gray catbird	Mimidae	Passeriformes	stable	Neotropic	[S60]
Northern black swift	Apodidae	Apodiformes	decreasing	Neotropic	[S61]
Osprey	Pandionidae	Accipitriformes	increasing	Neotropic	[S62–S65]

Ovenbird	Parulidae	Passeriformes	stable	Neotropic	[S66,S67]
Purple martin	Hirundinidae	Passeriformes	stable	Neotropic	[S68]
Red-eyed vireo	Vireonidae	Passeriformes	increasing	Neotropic	[S69]
Scissor-tailed flycatcher	Tyrannidae	Passeriformes	decreasing	Neotropic	[S59]
Veery	Turdidae	Passeriformes	decreasing	Neotropic	[S70,S71]
Western kingbird	Tyrannidae	Passeriformes	increasing	Neotropic	[S59]
Wood thrush	Turdidae	Passeriformes	decreasing	Neotropic	[S72]

217
218
219
220
221

222

223

224

225 Table S2. Full and best model summary table. w.spread = non-breeding mean spread; mig.distance =
226 distance of migration; system = Neotropical or Palearctic flyway; zone = location of non-breeding
227 ground in either Central or South America, or Africa approximately above and below the equator;
228 b.lon.rel = scaled mean breeding longitude; b.lat.rel = scaled mean breeding latitude.
229

	Parameter estimate														
		<i>Log</i>	<i>mig.</i>					<i>mig.</i>							
Model	<i>Intercept</i>	<i>(w.spread)</i>	<i>distance</i>	<i>system</i>	<i>zone</i>	<i>b.lon.rel</i>	<i>b.lat.rel</i>	<i>x system</i>	<i>w.spread</i>	<i>x system</i>	k	AICc	Delta	R ² m	R ² c
1	-0.50	-0.26	-0.026	0.46				-0.42	0.80		9	100.1	0	0.30	0.35
2	-0.47	-0.27	-0.0056	0.49	-0.082	0.00023	0.000097	-0.42	0.81		14	106.5	6.4	0.29	0.34

230
231
232

233 **Supplemental References**

- 234 S1. Finch, T., Butler, S., Franco, A., and Cresswell, W. (2017). Low migratory connectivity is
235 common in long-distance migrant birds. *J. Anim. Ecol.* **38**, 42–49.
- 236 S2. BirdLife International (2017). IUCN Red List for Birds. Available at: <http://www.birdlife.org>
237 [Accessed October 2, 2017].
- 238 S3. Liechti, F., Witvliet, W., Weber, R., and Bachler, E. (2013). First evidence of a 200-day non-
239 stop flight in a bird. *Nat. Commun.* **4**, 2554.
- 240 S4. Salewski, V., Flade, M., Poluda, A., Kiljan, G., Liechti, F., Lisovski, S., and Hahn, S. (2013). An
241 unknown migration route of the “globally threatened” aquatic warbler revealed by geolocators.
242 *J. Ornithol.* **154**, 549–552.
- 243 S5. Liechti, F., Scandolara, C., Rubolini, D., Ambrosini, R., Korner-Nievergelt, F., Hahn, S.,
244 Lardelli, R., Romano, M., Caprioli, M., Romano, A., *et al.* (2015). Timing of migration and
245 residence areas during the non-breeding period of barn swallows *Hirundo rustica* in relation to
246 sex and population. *J. Avian Biol.* **46**, 254–265.
- 247 S6. Willemoes, M., Strandberg, R., Klaassen, R.H.G., Tøttrup, A.P., Vardanis, Y., Howey, P.W.,
248 Thorup, K., Wikelski, M., and Alerstam, T. (2014). Narrow-front loop migration in a population
249 of the common cuckoo *Cuculus canorus*, as revealed by satellite telemetry. *PLoS One* **9**, 1–
250 10.
- 251 S7. Hewson, C.M., Thorup, K., Pearce-Higgins, J.W., and Atkinson, P.W. (2016). Population
252 decline is linked to migration route in the Common Cuckoo. *Nat. Commun.* **7**, 12296.
- 253 S8. Hahn, S., Emmenegger, T., Lisovski, S., Amrhein, V., Zehndjiev, P., and Liechti, F. (2014).
254 Variable detours in long-distance migration across ecological barriers and their relation to
255 habitat availability at ground. *Ecol. Evol.* **4**, 4150–4160.
- 256 S9. Cresswell, B., and Edwards, D. (2012). Geolocators reveal wintering areas of European
257 Nightjar (*Caprimulgus europaeus*). *Bird Study* **60**, 77–86.
- 258 S10. Norevik, G., Åkesson, S., and Hedenström, A. (2017). Migration strategies and annual space-
259 use in an Afro-Palaeartic aerial insectivore – the European nightjar *Caprimulgus europaeus*.
260 *J. Avian Biol.* **48**, 738–747.

- 261 S11. Evens, R., Conway, G.J., Ian, G., Cresswell, B., Ed, F.R., Witters, N., and Beenaerts, N.
262 (2017). Migratory pathways, stopover zones and wintering destinations of Western Nightjars
263 *Caprimulgus europaeus*. Ibis. 159, 680–686.
- 264 S12. Kristensen, M.W., Tøttrup, A.P., and Thorup, K. (2013). Migration of the Common Redstart
265 (*Phoenicurus phoenicurus*). Auk 130, 258–264.
- 266 S13. Åkesson, S., Klaassen, R., Holmgren, J., Fox, J.W., and Hedenström, A. (2012). Migration
267 routes and strategies in a highly aerial migrant, the common swift *Apus apus*, revealed by
268 light-level geolocators. PLoS One 7, 1–9.
- 269 S14. Klaassen, R., Klaassen, H., Berghuis, A., Berghuis, M., Schreven, K., Van Der Horst, Y.,
270 Verkade, H., and Kearsley, L. (2014). Trekroutes en overwinteringsgebieden van Nederlandse
271 Gierzwaluwen ontrafeld met geolocators. Limosa 87, 173–181.
- 272 S15. Åkesson, S., Bianco, G., and Hedenstrom, A. (2016). Negotiating an ecological barrier:
273 crossing the Sahara in relation to winds by common swifts. Philos. Trans. R. Soc. London B
274 Biol. Sci. 371, 20150393.
- 275 S16. Wellbrock, A.H.J., Bauch, C., Rozman, J., and Witte, K. (2016). “Same procedure as last
276 year?” – Repeatedly tracked swifts show individual consistency in migration pattern in
277 successive years. J. Avian Biol. 48, 897–903.
- 278 S17. Xenophontos, M., Blackburn, E., and Cresswell, W. (2017). Cyprus wheatears *Oenanthe*
279 *cypriaca* likely reach sub-Saharan African wintering grounds in a single migratory flight. J.
280 Avian Biol. 48, 1–7.
- 281 S18. García-Ripollés, C., López-López, P., and Urios, V. (2010). First description of migration and
282 wintering of adult Egyptian Vultures *Neophron percnopterus* tracked by GPS satellite
283 telemetry. Bird Study 57, 261–265.
- 284 S19. López-López, P., García-Ripollés, C., and Urios, V. (2014). Individual repeatability in timing
285 and spatial flexibility of migration routes of trans-Saharan migratory raptors. Curr. Zool. 60,
286 642–652.
- 287 S20. Gschweng, M., Kalko, E.K. V., Querner, U., Fiedler, W., and Berthold, P. (2008). All across
288 Africa: Highly individual migration routes of Eleonora’s falcon. Proc. R. Soc. B 275, 2887–

- 289 2896.
- 290 S21. Kassara, C., Fric, J., Gschweng, M., and Sfenthourakis, S. (2012). Complementing the puzzle
291 of Eleonora's Falcon (*Falco eleonorae*) migration: New evidence from an eastern colony in the
292 Aegean Sea. *J. Ornithol.* 153, 839–848.
- 293 S22. Mellone, U., López-López, P., Limiñana, R., Piasevoli, G., and Urios, V. (2013). The trans-
294 equatorial loop migration system of Eleonora's falcon: Differences in migration patterns
295 between age classes, regions and seasons. *J. Avian Biol.* 44, 417–426.
- 296 S23. Bächler, E., Hahn, S., Schaub, M., Arlettaz, R., Jenni, L., Fox, J.W., Afanasyev, V., and
297 Liechti, F. (2010). Year-round tracking of small trans-Saharan migrants using light-level
298 geolocators. *PLoS One* 5, 3–6.
- 299 S24. van Wijk, R.E., Bauer, S., and Schaub, M. (2016). Repeatability of individual migration routes,
300 wintering sites and timing in a long-distance migrant bird. *Ecol. Evol.* 6, 8679–8685.
- 301 S25. Finch, T., Saunders, P., Avilés, J.M., Bermejo, A., Catry, I., de la Puente, J., Emmenegger, T.,
302 Mardegá, I., Mayet, P., Parejo, D., *et al.* (2015). A pan-European, multipopulation assessment
303 of migratory connectivity in a near-threatened migrant bird. *Divers. Distrib.* 21, 1051–1062.
- 304 S26. Lemke, H.W., Tarka, M., Klaassen, R.H.G., Åkesson, M., Bensch, S., Hasselquist, D., and
305 Hansson, B. (2013). Annual cycle and migration strategies of a trans-saharan migratory
306 songbird: A geolocator study in the great reed warbler. *PLoS One* 8, 1–10.
- 307 S27. Koleček, J., Prochazka, P., El-Arabany, N., Tarka, M., Ilieva, M., Hahn, S., Honza, M., de la
308 Puente, J., Bermejo, A., Gursø, A., *et al.* (2016). Cross-continental migratory connectivity and
309 spatiotemporal migratory patterns in the great reed warbler. *J. Avian Biol.* 47, 756–767.
- 310 S28. Meyburg, B.U., Howey, P.W., Meyburg, C., and Fiuczynski, K.D. (2011). Two complete
311 migration cycles of an adult Hobby tracked by satellite. *Br. Birds* 104, 2–15.
- 312 S29. Strandberg, R., Klaassen, R.H., Hake, M., Olofsson, P., and Alerstam, T. (2009). Converging
313 migration routes of Eurasian hobbies *Falco subbuteo* crossing the African equatorial rain
314 forest. *Proc. R. Soc. B* 276, 727–733.
- 315 S30. Hake, M., Kjellen, N., and Alerstam, T. (2003). Age-dependent migration strategy in honey
316 buzzards *Pernis apivorus* tracked by satellite. *Oikos* 103, 385–396.

- 317 S31. Szép, T., Liechti, F., Nagy, K., Nagy, Z., and Hahn, S. (2017). Discovering the migration and
318 non-breeding areas of sand martins and house martins breeding in the Pannonian basin
319 (central-eastern Europe). *J. Avian Biol.* 48, 114–122.
- 320 S32. Catry, I., Dias, M.P., Catry, T., Afanasyev, V., Fox, J., Franco, A.M.A., and Sutherland, W.J.
321 (2010). Individual variation in migratory movements and winter behaviour of Iberian Lesser
322 Kestrels *Falco naumanni* revealed by geolocators. *Ibis.* 153, 154–164.
- 323 S33. Limiñana, R., Romero, M., Mellone, U., and Urios, V. (2012). Mapping the migratory routes
324 and wintering areas of Lesser Kestrels *Falco naumanni*: New insights from satellite telemetry.
325 *Ibis.* 154, 389–399.
- 326 S34. Meyburg, B.-U., Scheller, W., and Meyburg, C. (1995). Migration and wintering of the Lesser
327 spotted eagle (*Aquila pomarina*): a study by means of satellite telemetry. *J. Ornithol.* 136, 401–
328 422.
- 329 S35. Strandberg, R., Klaassen, R.H.G., Hake, M., Olofsson, P., Thorup, K., and Alerstam, T. (2008).
330 Complex timing of Marsh Harrier *Circus aeruginosus* migration due to pre- and post-migratory
331 movements. *Ardea* 96, 159–171.
- 332 S36. Trierweiler, C., Koks, B.J., Drent, R.H., Exo, K.M., Komdeur, J., Dijkstra, C., and Bairlein, F.
333 (2007). Satellite tracking of two Montagu's Harriers (*Circus pygargus*): Dual pathways during
334 autumn migration. *J. Ornithol.* 148, 513–516.
- 335 S37. Trierweiler, C., Drent, R.H., Komdeur, J., Exo, K.M., Bairlein, F., and Koks, B.J. (2008). De
336 jaarcyclus van de Grauwe Kiekendief: Een leven gedreven door woelnumuizen sprinkhanen.
337 *Limosa* 81, 107–115.
- 338 S38. Trierweiler, C., Klaassen, R.H.G., Drent, R.H., Exo, K.-M., Komdeur, J., Bairlein, F., and Koks,
339 B.J. (2014). Migratory connectivity and population-specific migration routes in a long-distance
340 migratory bird. *Proc. R. Soc. B* 281, 20132897.
- 341 S39. Schmaljohann, H., Buchmann, M., Fox, J.W., and Bairlein, F. (2012). Tracking migration
342 routes and the annual cycle of a trans-Sahara songbird migrant. *Behav. Ecol. Sociobiol.* 66,
343 915–922.
- 344 S40. Van Oosten, H.H., Versluijs, R., and Van Wijk, R. (2014). Migration routes and wintering areas

of two Dutch Northern Wheatears *Oenanthe oenanthe* in the Sahel. *Limosa* 87, 168–172.

- S41. Selstam, G., Sondell, J.A.N., and Olsson, P. (2015). Wintering area and migration routes for Ortolan Buntings *Emberiza hortulana* from Sweden determined with light-geologgers. *Ornis Svecica* 25, 3–14.
- S42. Jiguet, F., Arlettaz, R., Belik, V., Bernardy, P., Copete, J.L., Czajkowski, M.A., Dale, S., Dombrowski, V., Elts, J., Ferrand, Y., *et al.* (2016). Migration strategy of the ortolan bunting final report of the scientific committee. Museum Natl. D'Histoire Nat.
- S43. Terraube, J., Mougeot, F., Cornulier, T., Verma, A., Gavrilov, A., and Arroyo, B. (2012). Broad wintering range and intercontinental migratory divide within a core population of the near-threatened pallid harrier. *Divers. Distrib.* 18, 401–409.
- S44. Ouwehand, J., Ahola, M.P., Ausems, A.N.M.A., Bridge, E.S., Burgess, M., Hahn, S., Hewson, C.M., Klaassen, R.H.G., Laaksonen, T., Lampe, H.M., *et al.* (2016). Light-level geolocators reveal migratory connectivity in European populations of pied flycatchers *Ficedula hypoleuca*. *J. Avian Biol.* 47, 69–83.
- S45. Tøttrup, A.P., Klaassen, R.H.G., Strandberg, R., Thorup, K., Kristensen, M.W., Jørgensen, P.S., Fox, J., Afanasyev, V., Rahbek, C., and Alerstam, T. (2012). The annual cycle of a trans-equatorial Eurasian-African passerine migrant: different spatio-temporal strategies for autumn and spring migration. *Proc. R. Soc. B* 279, 1008–16.
- S46. Tøttrup, A.P., Pedersen, L., Onrubia, A., Klaassen, R.H.G., and Thorup, K. (2017). Migration of red-backed shrikes from the Iberian Peninsula: optimal or sub-optimal detour? *J. Avian Biol.* 48, 149–154.
- S47. Briedis, M., Traff, J., Hahn, S., Ilieva, M., Kral, M., Peev, S., and Adamik, P. (2016). Year-round spatiotemporal distribution of the enigmatic Semi-collared Flycatcher *Ficedula semitorquata*. *J. Ornithol.* 157, 895–900.
- S48. Meyburg, B.U., Meyburg, C., and Pacteau, C. (1996). Migration automnale d'un circaète Jean-le-Blanc *Circaetus gallicus* suivi par satellite. *Alauda* 64, 339–344.
- S49. Meyburg, B.U., Meyburg, C., and Barbraud, J.C. (1998). Migration strategies of an adult short-

- 373 toed eagle *Circaetus gallicus* tracked by satellite. *Alauda* 66, 39–48.
- 374 S50. Briedis, M., Beran, V., Hahn, S., and Adamík, P. (2016). Annual cycle and migration strategies
375 of a habitat specialist, the Tawny Pipit *Anthus campestris*, revealed by geolocators. *J. Ornithol.*
376 157, 619–626.
- 377 S51. Stach, R., Jakobsson, S., Kullberg, C., and Fransson, T. (2012). Geolocators reveal three
378 consecutive wintering areas in the thrush nightingale. *Anim. Migr.* 338, 1–7.
- 379 S52. Eraud, C., Riviere, M., Lormee, H., Fox, J.W., Ducamp, J.J., and Boutin, J.M. (2013). Migration
380 routes and staging areas of trans-Saharan turtle doves appraised from light-level geolocators.
381 *PLoS One* 8.
- 382 S53. Lormée, H., Boutin, J.M., Pinaud, D., Bidault, H., and Eraud, C. (2016). Turtle Dove
383 *Streptopelia turtur* migration routes and wintering areas revealed using satellite telemetry. *Bird*
384 Study 63, 425–429.
- 385 S54. Lerche-jørgensen, M., Willemoes, M., Tøttrup, A.P., Snell, K.R.S., and Thorup, K. (2017). No
386 apparent gain from continuing migration for more than 3000 kilometres: willow warblers
387 breeding in Denmark winter across the entire northern Savannah as revealed by geolocators.
388 *Movement* 5, 17.
- 389 S55. Pillar, A.G., Marra, P.P., Flood, N.J., and Reudink, M.W. (2016). Molt migration in Bullock's
390 orioles (*Icterus bullockii*) confirmed by geolocators and stable isotope analysis. *J. Ornithol.*
391 157, 265–275.
- 392 S56. DeLuca, W. V., Woodworth, B.K., Rimmer, C.C., Marra, P.P., Taylor, P.D., McFarland, K.P.,
393 Mackenzie, S.A., and Norris, D.R. (2015). Transoceanic migration by a 12 g songbird. *Biol.*
394 Lett. 11, 20141045.
- 395 S57. Renfrew, R.B., Kim, D., Perlut, N., Smith, J., Fox, J., and Marra, P.P. (2013). Phenological
396 matching across hemispheres in a long-distance migratory bird. *Divers. Distrib.* 19, 1008–
397 1019.
- 398 S58. Haines, A.M., Mcgrady, M.J., Martell, M.S., Dayton, B.J., Henke, M.B., and Seegar, W.S.
399 (2003). Migration routes and wintering locations of broad-winged hawks tracked by satellite
400 telemetry. *Wilson Bull.* 115, 166–169.

- 401 S59. Jahn, A.E., Cueto, V.R., Fox, J.W., Husak, M.S., Kim, D.H., Landoll, D. V., Ledezma, J.P.,
402 LePage, H.K., Levey, D.J., Murphy, M.T., *et al.* (2013). Migration timing and wintering areas of
403 three species of flycatchers (*Tyrannus*) breeding in the Great Plains of North America. *Auk*
404 130, 247–257.
- 405 S60. Ryder, T.B., Fox, J.W., and Marra, P.P. (2011). Estimating migratory connectivity of Gray
406 catbirds (*Dumetella carolinensis*) using geolocator and mark–recapture data. *Auk* 128, 810–
407 810.
- 408 S61. Beason, J.P., Gunn, C., Potter, K.M., Sparks, R.A., and Fox, J.W. (2012). The northern black
409 swift: Migration path and wintering area revealed. *Wilson J. Ornithol.* 124, 1–8.
- 410 S62. Martell, M.S., Henny, C.J., Nye, P.E., and Solensky, M.J. (2001). Fall migration routes, timing,
411 and wintering sites of North American ospreys as determined by satellite telemetry. *Condor*
412 103, 715–724.
- 413 S63. Martell, M.S., McMillan, M.A., Solensky, M.J., and Mealey, B.K. (2004). Partial migration and
414 wintering use of Florida by Ospreys. *J. Raptor Res.* 38, 55–61.
- 415 S64. Elliott, J.E., Morrissey, C.A., Henny, C.J., Inzunza, E.R., and Shaw, P. (2007). Satellite
416 telemetry and prey sampling reveal contaminant sources to pacific northwest ospreys. *Ecol.*
417 *Appl.* 17, 1223–1233.
- 418 S65. Bedrosian, B., Cain, S., Wolff, S., and Craighead, D. (2015). Migratory pathways, timing, and
419 home ranges of southern greater yellowstone osprey. *J. Raptor Res.* 49, 325–332.
- 420 S66. Hallworth, M.T., and Marra, P.P. (2015). Miniaturized GPS Tags Identify Non-breeding
421 Territories of a Small Breeding Migratory Songbird. *Sci. Rep.* 5, 1–6.
- 422 S67. Hallworth, M.T., Sillett, T.S., Van Wilgenburg, S.L., Hobson, K.A., and Marra, P.P. (2015).
423 Migratory connectivity of a Neotropical migratory songbird revealed by archival light-level
424 geolocators. *Ecol. Appl.* 25, 336–347.
- 425 S68. Fraser, K.C., Stutchbury, B.J.M., Silverio, C., Kramer, P.M., Barrow, J., Newstead, D., Mickle,
426 N., Cousens, B.F., Lee, J.C., Morrison, D.M., *et al.* (2012). Continent-wide tracking to
427 determine migratory connectivity and tropical habitat associations of a declining aerial
428 insectivore. *Proc. R. Soc. B* 279, 4901–4906.

- 429 S69. Callo, P.A., Morton, E.S., and Stutchbury, B.J.M. (2013). Prolonged spring migration in the
430 Red-eyed Vireo (*Vireo olivaceus*). *Auk* 130, 240–246.
- 431 S70. Hobson, K.A., and Kardynal, K.J. (2015). Western Veeries use an eastern shortest-distance
432 pathway: New insights to migration routes and phenology using light-level geolocators. *Auk*
433 132, 540–550.
- 434 S71. Heckscher, C.M., Halley, M.R., and Stampul, P.M. (2015). Intratropical migration of a nearctic-
435 neotropical migratory songbird (*Catharus fuscescens*) in south america with implications for
436 migration theory. *J. Trop. Ecol.* 31, 285–289.
- 437 S72. Stanley, C.Q., Mckinnon, E.A., Fraser, K.C., Macpherson, M.P., Casbourn, G., Friesen, L.,
438 Marra, P.P., Studds, C., Ryder, T.B., Diggs, N.E., *et al.* (2015). Connectivity of wood thrush
439 breeding, wintering, and migration sites based on range-wide tracking. *Conserv. Biol.* 29, 164–
440 174.
- 441
- 442